Fairview Community Primary School

• Phone parent/emergency contact





Child Name:	
Pate of Birth:	Form completed date:
Allergic to:	Review Date:
Mild/moderate reaction: • Swollen lips, face or eyes	Emergency Contacts:
Itchy/tingling mouthHives or itchy skin rash	Name:
Abdominal pain or vomitingSudden change in behaviour	Tel:
Action to take: • Stay with the child, call for help if necessary	Name:
 Locate adrenaline autoinjector(s) (EpiPen) Give antihistamine: 	Tel:
(Name of medication) (If vomited, repeat dose)	

How to give EpiPen

Watch for signs of ANAPHYLAXIS

Anaphylaxis may occur without skin symptoms ALWAYS consider anaphylaxis in someone with known food allergy who has SUDDEN BREATHING DIFFICULTY.

A AIRWAY

B BREATHING

C CONSCIOUSNESS

Persistent cough **Hoarse voice** Difficulty swallowing Wheeze or Swollen tongue

Difficult or noisy breathing persistent cough

Persistent dizziness Pale or floppy Suddenly sleepy Collapse/unconscious


IF ANY ONE (OR MORE) OF THESE SIGNS ABOVE ARE PRESENT:

- Lie child flat with legs raised (if breathing is difficult, allow child
- Use EpiPen without delay
- Dial 999 for ambulance and say ANAPHYLAXIS ***IF IN DOUBT, GIVE ADRENALINE ***

AFTER GIVING ADRENALINE:

- Stay with the child until ambulance arrives, do NOT stand child up
- Commence CPR if there are no signs of life 2.
- Phone parent/emergency contact
- If no improvement after 5 minutes, give a further adrenaline dose using a second EpiPen if available.

PULL OFF BLUE SAFETY CAP and grasp EpiPen. Remember "blue to sky, orange to the thigh".



PUSH DOWN HARD until a click is heard or felt and hold in place for 3 seconds. Remove EpiPen.







Additional instructions:

If wheezy, GIVE ADRENALINE FIRST, then asthma reliever (blue puffer) via spacer